

547
HER

Cey. R.

STRUCTURAL STUDIES OF A POLYSACCHARIDE FROM
LITSEA GLUTINOSA AND
THE SYNTHESIS OF SOME ACRIDONES AND COUMARINS

A THESIS SUBMITTED BY

H. M. T. BANDARA HERATH

IN PARTIAL FULFILMENT OF THE
REQUIREMENTS FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

OF THE


UNIVERSITY OF PERADENIYA


SRI LANKA

November 1991

Research Laboratory
Department of Chemistry

435565





ABSTRACT

This thesis consists of three parts.

Part-1 describes the structural studies of an arabinoxylan isolated from *Litsea glutinosa* (Lour).

Arabinose and the xylose were the major neutral sugars present in the water soluble polysaccharide isolated from the mucilaginous bark of *Litsea glutinosa* (Lauraceae). The molar ratio of D-xylose : L-arabinose was 1.0 : 3.5. Methylation analysis, partial hydrolysis and ^1H , ^1H - ^1H 2D-COSY, and ^{13}C -nmr spectroscopy indicated a backbone of (1-4)-linked β -D-xylopyranosyl residues substituted at both positions 2 and 3 with side chains composed of either single or (1-3)-linked arabino- furanosyl residues. Both α -L- and β -L- arabinofuranosyl residues were present. Side chains composed of two β -L-arabinofuranosyl residues are probably attached at O-2 of some xylosyl residues.

Part-2 of the thesis discusses the total synthesis of twenty acridone alkaloids, eleven of which are new acridones.

1,3-Dihydroxy-9-acridone (31) was synthesised from methyl anthranilate and phloroglucinol. Selective methylations of compound (31), gave 1-hydroxy-3-

methoxy-10-methyl-9-acridone (32), 1,3-dimethoxy-10-methyl-9-acridone (33) and 1,3-dihydroxy-10-methyl-9-acridone (34). The iodination of (31) afforded a new acridone, 1,3-dihydroxy-2,4-diiodo-9-acridone (35) while compounds (32), (33) and (34) gave 1-hydroxy-2-iodo-3-methoxy-10-methyl-9-acridone (36), 2-iodo-1,3-dimethoxy-10-methyl-9-acridone (37) and 1,3-dihydroxy-2-iodo-10-methyl-9-acridone (38) respectively. Methylation of compound (35) gave another new acridone, 2,4-diiodo-1,3-dimethoxy-10-methyl-9-acridone (39). The palladium catalysed Heck condensation reaction of compounds (36), (38) and (39) with 2-methyl-3-butene-2-ol afforded three new acridone alkaloids, 1-hydroxy-2-(3-hydroxy-3-methyl-1-butenyl)-3-methoxy-10-methyl-9-acridone (40), 1,3-dimethoxy-2-(3-hydroxy-3-methyl-1-butenyl)-10-methyl-9-acridone (42) and 2,4-bis(3-hydroxy-3-methyl-1-butenyl)-1,3-dimethoxy-10-methyl-9-acridone (43) respectively, while compound (37) afforded a naturally occurring acridone, isonoracronycine (41). Methylation of (41) gave isoacronycine (44).

Methyl-6-amino-2,3-dimethoxybenzoate (51) which was synthesised from 6-formyl-2,3-dimethoxybenzoic acid (45), was treated with phloroglucinol to give two new acridone alkaloids, 1,3,8-trihydroxy-7-methoxy-9-acridone (52) and 1,3-dihydroxy-7,8-dimethoxy-9-acridone (53). The selective methylation of (52) and

(53) gave another four new acridones, 1,8-dihydroxy-3,7-dimethoxy-9-acridone (54), 1,8-dihydroxy-3,7-dimethoxy-10-methyl-9-acridone (55), 1-hydroxy-3,7,8-trimethoxy-10-methyl-9-acridone and 1,3,7,8-tetramethoxy-10-methyl-9-acridone (57).

Part-3 of the thesis deals with the total synthesis of (*E*)-suberenol (3) from 4-bromoresorcinol and describes the use of (*E*)-suberenol as a precursor for the synthesis of related coumarin derivatives.

The synthesis of three naturally occurring coumarins, cyclobisuberodiene (9), dihydrosuberenol (11) and ethylsuberenol (12) and three new coumarins, (*E*)-suberodiene (8), (*E*)-suberenene (10) and methylsuberenol (13), a new coumarin dimer (14) and three new acridone-coumarin dimers (Acrimarins), 1,3-dihydroxy-2-[1-7-(methoxy-2-oxo-2*H*-chromen-6-yl)-3-methyl-2-butenyl]-10-methyl-9-acridone (15), 1-hydroxy-3-methoxy-2-[1-7-(methoxy-2-oxo-2*H*-chromen-6-yl)-3-methyl-2-butenyl]-10-methyl-9-acridone (16) and 1,3-dihydroxy-2-[1-7-(methoxy-2-oxo-2*H*-chromen-6-yl)-3-methyl-2-butenyl]-9-acridone (17) are described in this part of the thesis.