

OPTIMIZATION OF TOTAL RNA EXTRACTION FROM HUMAN URINARY SEDIMENT

S. Saseevan^{1,2}, D.N. Magana-Arachchi^{1*} and S. Rajapakse^{2,3}

¹National Institute of Fundamental Studies, Kandy, Sri Lanka

²Postgraduate Institute of Science, University of Peradeniya, Peradeniya, Sri Lanka

³Department of Molecular Biology and Biotechnology, Faculty of Science, University of Peradeniya, Peradeniya, Sri Lanka

*dhammika.ma@nifs.ac.lk

Urine is the best choice to identify biomarkers for metabolic and renal disorders because it is readily available, and samples can be obtained non-invasively from patients. However, RNA isolation from voided urine is challenging due to the presence of RNases and cell scarcity. This study aims to optimize a protocol for RNA extraction from urine samples in gene expression studies. Twenty urine samples were collected from healthy controls (HC) ($n = 11$; 49 ± 5 years) and chronic kidney disease (CKD) patients ($n = 9$; 62 ± 3 years) and were centrifuged at 3,000 g for 30 min at 4 °C. Then, 500 μ L of the lysis buffer was added to the pellet, vortexed and kept on ice for 5 min. Next, 100 μ L of sodium acetate (pH = 4.0) and 500 μ L of water-saturated phenol were added and mixed well. After that, 200 μ L of chloroform: isoamyl alcohol (49:1) was added, vortexed and centrifuged. An equal volume of cold isopropanol was added to the aqueous phase and incubated at -20 °C for 1 h to precipitate RNA. The pellet was washed with 75% ethanol, air dried, and resuspended with 12 μ L nuclease-free water. Finally, the RNA was quantified and reverse transcribed into cDNA to be used in RT-qPCR. Mean urine volume was 82.5 ± 41.9 mL. Serum creatinine and estimated glomerular filtration rate of CKD patients were 3.0 ± 0.2 mg dL⁻¹ and 19.2 ± 4.8 mL min⁻¹ 1.73 m⁻², respectively. The total yield of RNA from CKD and HC samples were 873 ± 523 ng and 735 ± 291 ng, respectively, and a statistically significant difference was not observed between the two study groups ($p > 0.05$). The $\beta 2$ microglobulin gene could be successfully amplified using samples even with a low cDNA concentration (0.625 ng). This modified phenol-chloroform based urinary RNA isolation method is less expensive, does not require RNA clean-up kits and provides a higher yield of RNA with less inhibition which is sufficient for downstream applications than column-based techniques.

Keywords: Chronic kidney disease, RNA isolation, Urinary sediment