

C  
516.78  
KUM

(1)

**A COMBINED TEST FOR THE MEDIAN AND THE EFFECTS OF  
FIVE NORMALITY TESTS ON THE COMBINED TEST**

A PROJECT REPORT PRESENTED BY

KUMBALATARA ARACHCHIGE CHESMI DILHAN KUMBALATARA

to the Board of Study in Statistics & Computer Science of the  
POSTGRADUATE INSTITUTE OF SCIENCE

*in partial fulfillment of the requirement  
for the award of the degree of*

MASTER OF SCIENCE IN APPLIED STATISTICS

of the



**UNIVERSITY OF PERADENIYA**

**SRI LANKA**

**2010**

**641577**

# A COMBINED TEST FOR THE MEDIAN AND THE EFFECTS OF FIVE NORMALITY TESTS ON THE COMBINED TEST

**K.A.C.D. Kumbalatara**

Postgraduate Institute of Science

University of Peradeniya

Peradeniya

Sri Lanka

Testing for the median ( $\nu$ ) of a random variable  $X$  is a basic problem in statistics. When the distribution of  $X$  is symmetric and the moments are finite, the mean ( $\mu$ ) and the median ( $\nu$ ) are the same. For such situations a test regarding the median is equivalent a test regarding the mean. In particular when the distribution of  $X$  is normal a  $t$ -test can be used as the test for the median. Otherwise a nonparametric test like Binomial Sign test can be used as the test for the median. Therefore we considered the following "Combined Median test" for  $H_0: \nu = \nu_0$  vs  $H_1: \nu \neq \nu_0$ , where  $\nu_0$  is a given value.

Step 1: Test for the normality of  $X$ .

Step 2: If the normality is not rejected, do the  $t$ -test for the median.

Step 3: otherwise, do the Binomial Sign test for the median.

There are several tests available for the normality of  $X$ . In this study we investigated the effects of five well known normality tests, namely, the Kolmogorov-Smirnov test, Anderson Darling test, Shapiro-Wilk test, D'Agostino's K-Squared test and Cramer-von Mises test on the above Combined Median test.

It was found that the significance level of the Combined Median test is closest to the nominal significance level when the Shapiro-Wilk test is used as the normality test in step 1. It was also found that the power of the Combined Median test does not depend on the normality test in step 1. The results were obtained using Monte Carlo studies with different distributions for  $X$  and different sample sizes. Based on these results, we recommend the Combined Median test with Shapiro-Wilk test for normality in step 1.